

In Vitro Effect of Activated Recombinant Factor VII (rFVIIa) on Coagulation Properties of Human Blood at Hypothermic Temperatures

Bijan S. Kheirabadi, PhD, Angel V. Delgado, PhD, Michael A. Dubick, PhD, Michael R. Scherer, MA, Chriselda G. Fedyk, MS, John B. Holcomb, MD, and Anthony E. Pusateri, PhD

Background: Recombinant activated factor VII (rFVIIa) is currently administered off-label to control diffuse coagulopathic bleeding of patients with traumatic injuries. These patients are often cold, acidotic, and coagulopathic upon arrival and each responds differently to rFVIIa therapy. This study investigated the effects of hypothermia on clotting and the potential benefit of rFVIIa administration on blood coagulation at different hypothermic temperatures.

Method: Citrated blood samples were collected from eight healthy volunteers (20–45 years old) and incubated at 37°C, 34°C, 31°C, and 28°C for 30 minutes. rFVIIa (1.26 µg/mL equivalent to 90 µg/kg in vivo dose) or vehicle solution (saline) was added to each blood sample, incubated (10 minutes), and analyzed at the

respective temperatures by standard coagulation tests and thrombelastography.

Results: The clot reaction time of blood samples, measured as prothrombin time, activated partial thromboplastin time, and R time (thrombelastography analysis), was significantly prolonged at 31°C or below compared with at 37°C. The clot formation rate (α angle, maximum clotting velocity [V_{max}]) was decreased at all cold temperatures. Maximum clot strength (maximum amplitude) was only affected (reduced) at 28°C. Addition of rFVIIa shortened the prothrombin time, activated partial thromboplastin time, and R times at every temperature, surpassing the normal (37°C) temperature values in 31°C and 34°C cold samples. Similarly, clot formation rate parameters (clotting time, α angle, V_{max}) were also

improved by rFVIIa addition and normothermic values were restored in 31°C and 34°C cold blood samples. rFVIIa did not affect maximum amplitude at any temperature.

Conclusions: Mild to moderate hypothermia delayed the initial clot reaction and reduced clot formation rate without affecting ultimate clot strength. FVIIa effectively compensated for the adverse effects of hypothermia except in severe cases. These results suggest that administration of FVIIa should be beneficial in enhancing hemostasis in hypothermic trauma patients without the need for prior correction of the patient's body temperature.

Key Words: Hypothermia, Recombinant factor VIIa, Coagulation, Thrombelastography (TEG), Human blood.

J Trauma. 2007;63:1079–1086.

Uncontrollable bleeding accounts for 39% of trauma-related deaths and is the leading cause of potentially preventable death in patients with major trauma.^{1,2} A major contributing factor to bleeding-related mortality is coagulopathy, particularly when it is associated with metabolic acidosis and hypothermia, often referred to as the “lethal triad”.^{3–5} Substantial increases in mortality of trauma patients have been associated with profound hypothermia. Patients with admission temperatures less than 35°C had significantly greater mortality (25.5% vs. 3.0%) than patients with temperatures $\geq 35^\circ\text{C}$.⁶

Trauma with hypothermia has been shown to have significantly worse prognosis than either trauma or hypothermia alone.^{7–9} Although isolated hypothermia (32°C) incidents led to 23% mortality, hypothermia in conjunction with severe trauma (Injury Severity Score, [ISS] ≥ 25) was associated with 100% mortality in patients with core temperature below 32°C.^{8,9} This critical temperature was identified as an ominous predictor of mortality independent of the ISS, presence of shock, or the volume of fluid resuscitation. The mortality of hypothermic patients with comparable ISS, shock, and fluid resuscitation was significantly higher than that of those who remained warm.^{8,9}

A major risk factor of hypothermia is inducing coagulopathy with abnormal bleeding. Hypothermia can cause coagulopathy over a range of temperatures in a temperature-dependent fashion.⁸ Several mechanisms for the effect of hypothermia have been proposed, including reduced platelet function, decreased activity of coagulation enzymes, and activation of fibrinolysis; however, the exact mechanism continues to be debated.¹⁰

Recombinant activated factor VII (rFVIIa) has been considered as a “universal hemostatic agent”.¹¹ It was originally developed for the treatment of bleeding complications in

Submitted for publication May 21, 2007.

Accepted for publication August 15, 2007.

Copyright © 2007 by Lippincott Williams & Wilkins

From the Division of Damage Control Resuscitation, US Army Institute of Surgical Research, Fort Sam Houston, Texas.

The opinions or assertions expressed herein are the private views of the authors and are not to be construed as official or as reflecting the views of the US Department of the Army or the US Department of Defense.

Address for reprints: Bijan S. Kheirabadi, PhD, US Army Institute of Surgical Research, 3400 Rawley E. Chambers Avenue, Fort Sam Houston, TX 78234-6315; email: bijan.kheirabadi@cen.amedd.army.mil.

DOI: 10.1097/TA.0b013e31815885f1

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 01 NOV 2007		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE In vitro effect of activated recombinant factor VII (rFVIIa) on coagulation properties of human blood at hypothermic temperatures				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Kheirabadi B. S., Delgado A. V., Dubick M. A., Scherer M. R., Fedyk C. G., Holcomb J. B., Pusateri A. E.,				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) United States Army Institute of Surgical Research, JBSA Fort Sam Houston, TX 78234				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 8	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

hemophiliac patients with inhibitors against factors VIII or IX¹² and has been approved by the Food and Drug Administration for these indications. Successful treatment with this drug has been widely reported in hemophiliacs as well as other patients with pre-existing hemostatic defects (i.e., factor VIIa deficiency, acquired von Willebrand disease, uremia, and liver disease) for controlling acute bleeding episodes during surgical procedures.^{13,14} The off-label use of rFVIIa in a bleeding trauma patient, without preexisting hemostatic disorders, was first described in 1999.¹⁵ Since then, the emergency use of rFVIIa as an adjunct therapy for surgical or trauma patients with massive hemorrhage and coagulopathy has become an important new treatment option.^{16–25} It has provided a safe and effective means for control of severe bleeding when all other measures have failed. In a double-blinded, randomized controlled trial, rFVIIa treatment reduced transfusion requirements and the incidence of multiple organ failure and acute respiratory distress syndrome in severe blunt-trauma patients without increasing thromboembolic adverse effects.²⁶ The treatment, however, did not change the patients' mortality at either 48 hours or at 30 days after treatment.²⁶ In a recent clinical study, it was reported that rFVIIa may be able to improve early survival of massively bleeding trauma patients.²⁷ It is also suggested that the correction of acidosis and thrombocytopenia may be important prerequisites for rFVIIa efficacy.²⁷ The high cost associated with this treatment and its questionable life-saving effect in trauma patients support the need for continued research to identify better conditions for improving the efficacy of this drug. The purpose of this study was to investigate the effects of hypothermia on coagulation parameters of normal human blood and determine the efficacy of rFVIIa to enhance coagulation at different hypothermic temperatures.

PATIENTS AND METHODS

Blood Sampling and Treatment

The Institutional Review Board of the US Army Institute of Surgical Research and Brooke Army Medical Center approved this study. After obtaining informed consent, blood was collected from eight adult healthy volunteers, (four men and four nonpregnant women), 20 years to 45 years old. None of the participants had received medication with acetylsalicylic acid or any other nonsteroidal anti-inflammatory drugs for the 7 days before blood sampling. A smooth cubital venipuncture was performed using 21-gauge needle and blood (~65 mL) was collected into citrated vacutainer tubes (0.109 mol/L = 3.2% sodium citrate), mixing one part of citrate with nine parts of blood. The first tube aspirated was discarded to minimize tissue thromboplastin contamination. Blood from the remaining tubes was pooled in a 250 mL Nalgene PETG bottle (Fisher Scientific, Pittsburgh, PA) and incubated at 37°C for 15 minutes. After initial equilibration, the blood was aliquoted (6.5 mL) into small tubes and placed in temperature-controlled water baths at 37°C, 34°C, 31°C, or 28°C for an additional 30 minutes. Next, blood specimens

were treated with rFVIIa or vehicle. rFVIIa (1.2 mg kit, NovoSeven, Novo Nordisk, Denmark) was reconstituted in sterile water (2.2 mL) and 13.6 μ L of the solution was added to each 6.5 mL blood sample. Normal saline (13.6 μ L) was added to control blood. The final concentration of rFVIIa in blood was 1.26 μ g/mL, equivalent to a bolus dose of 90 μ g/kg in patients. The calculated concentration was based on the assumptions that rFVIIa will distribute evenly in 5 L circulating blood volume of an average patient with 70 kg body weight. The blood samples were kept at their normal or hypothermic temperatures for an additional 10 minutes and then assayed for coagulation function by the standard clinical tests and by thrombelastograph (TEG) analysis at their respective temperatures.

Coagulation Tests

After treatment, portions of blood samples were centrifuged at high speed, and the plasma collected for the standard clotting assays. Prothrombin time (PT), activated partial thromboplastin time (aPTT), and fibrinogen concentration were measured in plasma using automated BCS Coagulation Analyzer (Dade Behring, Marburg, Germany). All of the tests were performed at the corresponding experimental temperatures (28°C–37°C) in duplicate. For PT measurement, the clotting was induced by the addition of recombinant human tissue factor and synthetic phospholipids (Innovin, Dade Behring, Marburg, Germany) to the plasma in a 2:1 volume ratio. For aPTT measurement, purified soy phosphatides in ellagic acid (Actin FS Activated PTT Reagent, Dade Behring, Marburg, Germany) was added to plasma samples in a 1:1 volume ratio and clotting time was measured. Fibrinogen in plasma was also determined based on a clotting assay in which a large excess of bovine thrombin (50 IU/mL) was added to plasma in a 2:1 volume ratio, and the clotting time was measured. The fibrinogen concentration, which largely depends on this clotting time, was estimated based on a standard curve generated by the machine. Calcium was added to overcome the citrate anticoagulant according to standard practice.

TEG Analysis

Coagulation of all blood samples was tested with a TEG Hemostasis Analyzer 5000 (Hemoscope, Niles, IL). The accurate measurement by the TEG machines was confirmed daily using quality control standards obtained from Hemoscope. Before testing, the machines were set at the selected experimental temperatures of the blood samples. Disposable cups and pins were then loaded and allowed to equilibrate to the selected temperatures for at least 15 minutes. Next, 10 μ L Innovin (diluted 1:500 with saline), 20 μ L of 0.2 mol/L CaCl_2 , and 4.3 μ L of Corn Trypsin Inhibitor (19 μ g/mL, Hematologic Technologies Inc., Essex Junction, VT), a contact activation inhibitor, were added to each cup and allowed to equilibrate for an additional 5 minutes. To start the reaction, 336 μ L of citrated blood was added to each cup and measurement

started. Blood samples were tested in triplicate and TEG measurements continued until 30 minutes after maximum amplitude (MA) was reached. The following variables were measured for each sample at the experimental temperatures: *reaction time* (*R*, min, the time that the initial fibrin formation is detected and signaled by a 2 mm rise of amplitude); *clotting time* (*K*, min, indicates the speed of clot formation and is the time from the beginning of clotting, *R* time, until a clot with a fixed firmness is formed, 20 mm amplitude rise); *angle* (α degree, represents the kinetics of clot development); and MA (mm, measures the maximum strength or firmness of the developed clot). The velocity of clot formation was also calculated as the first derivative of the TEG tracings and maximum clotting velocity (*V*_{max}, mm/min) and time to reach *V*_{max} (*t-V*_{max}, min) were determined for each sample as described previously.²⁸ Fibrinolysis was measured as percent clot lysis at 30 minutes after MA was reached. This is measured based on reduction of the area under the TEG tracing between the two time points.

Statistical Analysis

All statistical analyses were performed using the statistical program SAS, version 8.1 (SAS Institute Inc. 1999, Cary, NC). Data were examined for heterogeneity of variance and non-normality. These conditions were not detected in the data. The coagulation measurements such as PT, aPTT, fibrinogen and TEG parameters were first compared among temperatures within each treatment group using a mixed model of analysis of variance, allowing for temperatures as fixed effects and replicate subject as a random effect. If there was a significant temperature effect, Dunnett's test was used to compare the normal (37°C) temperature with hypothermic temperatures. The parameters were then compared between the two treatment groups (rFVIIa vs. saline) at each temperature using a paired *t* test. Data are expressed as means \pm SEM in the graphs. Statistical significance was assigned at a greater than 95% confidence level ($p < 0.05$).

RESULTS

Plasma Clotting Assays

PT, which examines the *in vivo* (extrinsic) coagulation process induced by tissue factor was significantly prolonged in cold blood samples measured at 31°C and 28°C compared with 37°C (Fig. 1, top). The addition of rFVIIa to blood accelerated the initial clotting reaction and shortened the PT at every tested temperature. As a result, the PT of all rFVIIa treated samples was shorter than the value of untreated controls at 37°C (Fig. 1, top).

aPTT, a measurement of contact activation clotting (intrinsic) pathway, was more sensitive to low temperatures than PT (Fig. 1, bottom). The aPTT was prolonged at each hypothermic temperature in a temperature-dependent manner. The addition of rFVIIa to blood samples accelerated the clotting process and reduced aPTT at every temperature tested. The shortening of aPTT with rFVIIa compensated for the cold




Fig. 1. Prothrombin time (PT) (top) and activated partial thromboplastin time (aPTT), (bottom) of saline or rFVIIa added human blood samples at normal or hypothermic temperatures. PT and aPTT were significantly prolonged in cold blood compared with 37°C samples. rFVIIa accelerated the clotting process and reduced PT and aPTT at every temperature. Data represent mean \pm SEM of triplicate measurement of samples from eight subjects. + $p < 0.05$, significantly different from the 37°C temperature in each group. * $p < 0.05$, ** $p < 0.01$, significantly different from saline (control) group.

temperature effects measured in 31°C and 34°C blood samples (no difference compared with 37°C controls).

Fibrinogen measurement was the least affected by cold temperature. This assay is also based on a clotting time measurement but it does not depend on endogenous thrombin generation. The fibrinogen measurement was reduced (15%) only at the lowest temperature (28°C) compared with 37°C. rFVIIa addition had no significant effect on these measurements at any experimental temperature.

TEG Measurements

Typical TEG traces and velocity graphs of analysis of blood collected from a single subject at different temperatures are shown in Figure 2. Each tracing represents the average data obtained from triplicate tests. As seen in the figures, the cold temperatures adversely affected all aspects of clot formation except the clot strength. The detailed analysis of the TEG data, including the effect of rFVIIa at different temperatures, is as follows:

The initial clotting reaction time (*R*) was prolonged significantly (~ 1 – 2 minutes) at 31°C and 28°C compared with 37°C (Fig. 3). The addition of rFVIIa sped up the initial




Fig. 2. Typical TEG traces (left) and calculated velocity profiles (right) of blood from a subject at different temperatures. Each TEG trace represents the average of triplicate measurement of a blood sample at specific temperature. Velocity graphs were constructed from the triplicate TEG data measured at each temperature.

reaction and decreased the *R* times by ~ 2 minutes in all the samples, irrespective of their temperatures. The clot propagation was more affected by the low temperatures than the initial reaction time. These changes were reflected as longer clotting time (*K*) and smaller clotting rate (α angle) of cold blood samples compared with normothermic controls (Fig. 4). Again, the addition of rFVIIa was effective at every temperature, reducing the clotting times by 0.6 minutes to 0.9 minutes and increasing the clotting rate by $\sim 8^\circ$ at every temperature tested. The calculated clotting velocity values (*V*_{max} and *t-V*_{max}) also indicated slower clotting kinetics under hypothermic conditions (Fig. 5). The decrease of *V*_{max} and increase of *t-V*_{max} were significant at 31°C and 28°C compared with normal temperature. The addition of rFVIIa resulted in an increase in *V*_{max} and a decrease in *t-V*_{max} in all of the samples at every temperature tested. The improvement achieved by adding rFVIIa compensated or surpassed the changes caused by hypothermic temperatures.




Fig. 3. Initial clotting reaction time (*R*) of saline or rFVIIa added human blood samples at normal or hypothermic temperatures. *R*-time was prolonged significantly at 31°C and 28°C, compared with 37°C, and was shortened by addition of rFVIIa at each specific temperature. See Figure 1 for notation of significance.

The MA, which represents clot strength and depends on fibrinogen and platelet interaction, was the least affected by low temperatures (Fig. 6). A significant decrease in MA was measured only at the lowest temperature (28°C) in both untreated and rFVIIa-treated samples. The addition of rFVIIa had no effect on clot strength at any temperature. With regard to fibrinolysis, only a small fraction ($\sim 0.2\%$) of the clot lysed 30 minute post-MA (LY30) at 37°C. This reaction was also affected by cold temperatures and reduced to 0% at 31°C and




Fig. 4. Clot formation time (*K*) (top) and clotting rate (α angle) (bottom) of saline or rFVIIa added human blood samples at normal or hypothermic temperatures. The *K*-time increased and α angle decreased progressively with the decreasing of temperatures. rFVIIa addition shorten *K* time and fasten the clotting rate at each specific temperature. See Figure 1 for notation of significance.




Fig. 5. Maximum clotting velocity (V_{max}) (top) and time to reach V_{max} ($t-V_{max}$) (bottom) values of human blood at different temperatures calculated from TEG measurements. The decrease of V_{max} was significant at all hypothermic temperatures and $t-V_{max}$ was prolonged at 31°C and 28°C compared with 37°C. rFVIIa increased V_{max} and shortened $t-V_{max}$ in all the samples at every temperature. See Figure 1 for notation of significance.

28°C. There was no difference in clot lysis of rFVIIa-treated blood versus untreated samples.

DISCUSSION

Hypothermia occurs when the body core temperature falls below 35°C (95°F).²⁹ It has profound effects on every system of the body including reducing metabolism and oxygen demand, which may be beneficial, and decreasing the rate of temperature-dependent enzymatic function.³⁰ For exam-




Fig. 6. Maximum clot strength (MA) of saline or rFVIIa added human blood samples at normal or hypothermic temperatures. MA was least affected by hypothermia, it only decreased slightly at severe cold temperature (28°C) compared with 37°C. rFVIIa addition had no effect on MA. See Figure 1 for notation of significance.

ple, it is well recognized that, in general, for every 10°C decrease in temperature the activity of an enzyme is reduced by approximately 50%.³¹ This study investigated the effects of mild (34°C), moderate (31°C) and severe (28°C) hypothermia²⁹ on the coagulation of human blood and the potential ability of rFVIIa to reverse the coagulopathy and restore normal clotting function in hypothermic blood. The low temperatures affected the kinetics of clot formation by slowing (1) the cascade reactions of coagulation factors leading up to the formation of initial fibrin strands, which was detected as prolongation of PT, aPTT, and R-time in TEG analysis; and (2) the build up and cross linking of fibrin (propagation of clot), as determined by K time and clotting velocity measurements (α angle and V_{max}) by TEG analysis. The observed effects were temperature-dependent and proportional to the severity of hypothermia. These findings were consistent with other studies in which blood samples were collected from hypothermic patients and analyzed by TEG at corrected temperatures.^{32–34} The prolongation of PT and aPTT of normal blood when measured at hypothermic temperatures has been reported elsewhere.^{35,36} In clinics, these laboratory tests are usually performed on plasma samples after warming to 37°C, which can falsely produce normal values and underestimate coagulopathy in hypothermic patients.^{36,37}

Our TEG data showed that clot strength (MA), which represents the contribution of platelets and fibrinogen and their interaction to form a hemostatic clot, was not influenced by moderate hypothermia. The fibrinogen measurements estimated indirectly by adding excessive amounts of thrombin to plasma samples were also unaffected by hypothermia. These results collectively suggest that the moderate hypothermic temperatures slow down clot formation by decreasing enzymatic activity and reducing the thrombin generation rate without affecting the final contribution of platelets or fibrinogen to form the clot. The velocity profiles of hypothermic blood (Fig. 4), which display patterns similar to endogenous thrombin potential of the samples,³⁸ also suggested temperature-dependent decreases in thrombin generation. This mechanism is supported by an *in vitro* study which has shown that the monocyte tissue factor activities, measured as factor Xa generation, as well as platelet thrombin generation were reduced by 38% and 26%, respectively, at 33°C.³⁹

Watts et al. reported significant alterations in both enzymatic clotting reaction and platelet function (MA) of blood collected from hypothermic trauma patients with temperature below 34°C.³⁴ The TEG analysis also detected an acute hypercoagulability in blood of patients whose temperatures were $\geq 34^\circ\text{C}$. Other studies found no significant change in MA of blood collected from hypothermic patients,^{32,33} consistent with the present *in vitro* findings. The differing results reported among studies may be explained based on the difference in the sources of blood samples. The data reported by Watts et al.³⁴ were obtained from blood samples collected from hypothermic patients with significant traumatic injuries

(ISS ≥ 9) that may have adversely affected fibrinogen status and platelet function. The TEG analysis of blood collected from animals with moderate hypothermia also showed no change in the clot strength as compared with blood from normothermic animals.^{40,41}

In the present study, the *in vitro* addition of rFVIIa to the samples at a concentration considered to be the equivalent to a 90 $\mu\text{g/kg}$ *in vivo* dose corrected the coagulopathic effect of mild and moderate hypothermia (34°C and 31°C), but had less effect on severely hypothermic blood (28°C). rFVIIa shortened PT and aPTT of hypothermic samples and brought these clotting times to the level of normothermic blood but had no effect on fibrinogen measurements. This lack of effect can be explained by the fact that fibrinogen determination does not directly measure the concentration of the fibrinogen molecules in plasma but rather estimates the protein level from the clotting time of the sample when an excess amount of thrombin (an activated factor) is added. Thus, fibrinogen determination is not influenced by hypothermia because the clotting reaction bypasses the slowed coagulation process and delayed thrombin generation caused by hypothermia. rFVIIa addition would also have no effect on fibrinogen determination by this technique because by adding exogenous thrombin in excess, the need for FVIIa activity to propagate coagulation and enhance thrombin generation is totally circumvented.

To analyze the TEG findings, a summary of the changes in TEG parameters induced by hypothermia and the improvements achieved by the addition of rFVIIa as compared with the values of untreated blood at normal temperature are presented in Table 1. The effect of rFVIIa was temperature-dependent, improving both R-time and Vmax with greater effect on R-time. rFVIIa was most effective when added to mildly hypothermic blood; it not only corrected for the hypothermia effect but produced a clotting rate that surpassed the normothermic blood by ~25%. The increased coagulation rate observed here, may suggest the risk of thrombosis in patients treated with rFVIIa. However, rFVIIa is not indicated for treating hypothermic patient, it is rather targeted to the patients with significant coagulopathy and associated hypothermia and acidosis who are unresponsive to standard ther-

apy. In these situations, the drug reverses the coagulopathy and restores normal clotting function without increasing the risk of thrombosis.^{24,25,42,43}

With regard to contribution of platelet and fibrinogen in forming strong clot, rFVIIa did not have any impact even when it was added to normothermic blood. The effect of rFVIIa on fibrinolysis was also insignificant. The lack of effect of rFVIIa on clot strength (MA) was also reported in another *in vitro* study⁴⁴ and in an *in vivo* study²⁸ in which swine were injected with increasing doses of rFVIIa (180–720 $\mu\text{g/kg}$) and their blood was analyzed by TEG. These observations appear to be inconsistent with the reported effect of rFVIIa on increasing hemostatic strength of clot in an uncontrolled arterial hemorrhage model.⁴⁵ In that study, pretreatment of pigs with rFVIIa increased the clot strength and the pressure at which rebleeding occurred after fluid resuscitation.⁴⁵ The addition of rFVIIa, factor X, and prothrombin to washed platelets and red cell media (plasma free blood) was also shown to increase platelet adhesion and activation on collagen- and fibrinogen-coated surfaces.⁴⁶ These findings suggest that rFVIIa-mediated thrombin generation may enhance the hemostatic strength of clot and its attachment to injured tissues. However, this hemostatic improvement cannot be measured by the TEG analysis of blood that evaluates only the cohesive strength and not the adhesive strength of clot to the tissues.

The mechanism by which therapeutic doses of rFVIIa induce initial thrombin generation is still a matter of debate.^{47,48} According to a cell-based model, rFVIIa at pharmacological concentrations binds to activated platelets, exposing negatively charged phospholipids, and activates factor X independently from tissue factor.⁴⁹ This pathway of rFVIIa activity was not affected by the reduced temperature (33°C), suggesting rFVIIa should be effective in enhancing hemostasis in hypothermic patients.⁵⁰ Indeed, the effectiveness of rFVIIa to correct coagulopathy and stop bleeding in trauma patients with hypothermia was reported in the guidelines for the use of rFVIIa, and elsewhere.^{51,52} Experimental studies also showed that administration of rFVIIa in hemodiluted pigs improved coagulation parameters and reduced bleeding from a severe liver injury in the presence of hypothermia.^{53,54}

Table 1 Changes in TEG Parameters of Hypothermic Blood Samples With or Without rFVIIa Compared With Those of Untreated Blood at Normal Temperature (37°C)

Parameter	Temperature (°C)	Without rFVIIa	<i>p</i>	With rFVIIa	<i>p</i>
R-time (min)	34	18% slower	NS*	27% faster	0.02
	31	31% slower	0.02	17% faster	0.06
	28	37% slower	<0.01	4% slower	NS
Vmax (mm/min)	34	14% slower	0.02	24% faster	0.06
	31	22% slower	<0.01	3% slower	NS
	28	38% slower	<0.01	20% slower	<0.01
MA (mm)	34	1% reduced	NS	0% change	NS
	31	2% reduced	NS	2% reduced	NS
	28	4% reduced	<0.05	3% reduced	0.03

*NS, not significant, defined as *p* > 0.1.

However, the decreases in the bleeding in these studies and the subsequent investigation⁵⁵ with a greater number of animals did not result in a reduction in mortality rate.

The limitation of this study is that the rFVIIa effect was assessed in an isolated in vitro condition on blood collected from healthy volunteers, apart from many other influencing factors (e.g. tissue injury, hemorrhage, and acidosis) that may affect rFVIIa responses in trauma patients. On the other hand, this isolation is also the strength of this in vitro study that allows exploring the direct effect of hypothermia and defining the effectiveness of rFVIIa under different hypothermic conditions apart from other confounding factors that makes the interpretation of the findings very difficult.

In summary, hypothermia at 34°C, 31°C, and 28°C affected blood coagulation function by delaying the initial reaction process and decreasing the clot formation rate without impacting the final clot strength. Hypothermia alone, therefore, can be the cause of increased bleeding in trauma patients who may have normal clotting values (PT and aPTT measured at 37°C). The addition of rFVIIa to hypothermic blood compensated for the decreases and restored or increased the clotting reaction process above normothermic values in the cold blood samples. These data suggest that administration of rFVIIa should be beneficial in promoting hemostasis in hypothermic trauma patients without the need to correct the patient's body temperature before the treatment.

REFERENCES

- Hoyt DB, Bulger EM, Knudson MM, et al. Death in the operating room: an analysis of a multi-center experience. *J Trauma*. 1994; 37:426–432.
- Sauaia A, Moore FA, Moore EE, et al. Epidemiology of trauma deaths: a reassessment. *J Trauma*. 1995;38:185–193.
- Ferrara A, MacArthur JD, Wright HK, Modlin IM, McMillen MA. Hypothermia and acidosis worsen coagulopathy in the patient requiring massive transfusion. *Am J Surg*. 1990;160:515–518.
- Cosgriff N, Moore EE, Sauaia A, Kenny-Moynihan M, Burch JM, Galloway B. Predicting life-threatening coagulopathy in the massively transfused trauma patient: hypothermia and acidosis revisited. *J Trauma*. 1997;42:857–861; discussion 861–862.
- Lynn M, Jeroukhimov I, Klein Y, Martinowitz U. Updates in the management of severe coagulopathy in trauma patients. *Intensive Care Med*. 2002;28(suppl 2):S241–S247.
- Martin RS, Kilgo PD, Miller PR, et al. Injury-associated hypothermia: an analysis of the 2004 National Trauma Data Bank. *Shock*. 2005;24:114–118.
- Danzl DF, Pozos RS, Auerbach PS, et al. Multicenter hypothermia survey. *Ann Emerg Med*. 1987;16:1042–1055.
- Jurkovich GJ, Greiser WB, Luteran A, Curreri PW. Hypothermia in trauma victims: an ominous predictor of survival. *J Trauma*. 1987;27:1019–1024.
- Peng RY, Bongard FS. Hypothermia in trauma patients. *J Am Coll Surg*. 1999;188:685–696.
- Kirkpatrick AW, Chun R, Brown R, Simons RK. Hypothermia and the trauma patient. *Can J Surg*. 1999;42:333–343.
- Hedner U. Recombinant activated factor VII as a universal haemostatic agent. *Blood Coagul Fibrinolysis*. 1998;9(suppl 1): S147–S152.
- Hedner U, Glazer S, Pingel K, et al. Successful use of recombinant factor VIIa in patient with severe haemophilia A during synovectomy. *Lancet*. 1988;2:1193.
- Hay CR, Negrier C, Ludlam CA. The treatment of bleeding in acquired haemophilia with recombinant factor VIIa: a multicentre study. *Thromb Haemost*. 1997;78:1463–1467.
- Shapiro AD, Gilchrist GS, Hoots WK, Cooper HA, Gastineau DA. Prospective, randomised trial of two doses of rFVIIa (NovoSeven) in haemophilia patients with inhibitors undergoing surgery. *Thromb Haemost*. 1998;80:773–778.
- Kenet G, Walden R, Eldad A, Martinowitz U. Treatment of traumatic bleeding with recombinant factor VIIa. *Lancet*. 1999; 354:1879.
- Eikelboom JW, Bird R, Blythe D, et al. Recombinant activated factor VII for the treatment of life-threatening haemorrhage. *Blood Coagul Fibrinolysis*. 2003;14:713–717.
- Gowers CJ, Parr MJ. Recombinant activated factor VIIa use in massive transfusion and coagulopathy unresponsive to conventional therapy. *Anaesth Intensive Care*. 2005;33:196–200.
- Khan AZ, Parry JM, Crowley WF, et al. Recombinant factor VIIa for the treatment of severe postoperative and traumatic hemorrhage. *Am J Surg*. 2005;189:331–334.
- Kulkarni R, Daneshmand A, Guertin S, et al. Successful use of activated recombinant factor VII in traumatic liver injuries in children. *J Trauma*. 2004;56:1348–1352.
- Martinowitz U, Kenet G, Segal E, et al. Recombinant activated factor VII for adjunctive hemorrhage control in trauma. *J Trauma*. 2001;51:431–438; discussion 438–439.
- Mayo A, Misgav M, Kluger Y, et al. Recombinant activated factor VII (NovoSeven): addition to replacement therapy in acute, uncontrolled and life-threatening bleeding. *Vox Sang*. 2004;87:34–40.
- O'Neill PA, Bluth M, Gloster ES, et al. Successful use of recombinant activated factor VII for trauma-associated hemorrhage in a patient without preexisting coagulopathy. *J Trauma*. 2002; 52:400–405.
- Udy A, Vaghela M, Lawton G, Sigston P. The use of recombinant activated factor VII in the control of hemorrhage following blunt pelvic trauma. *Anaesthesia*. 2005;60:613–616.
- Dutton RP, McCunn M, Hyder M, et al. Factor VIIa for correction of traumatic coagulopathy. *J Trauma*. 2004;57:709–718; discussion 718–719.
- Haas T, Innerhofer P, Kuhbacher G, Fries D. Successful reversal of deleterious coagulopathy by recombinant factor VIIa. *Anesth Analg*. 2005;100:54–58.
- Boffard KD, Riou B, Warren B, et al. Recombinant factor VIIa as adjunctive therapy for bleeding control in severely injured trauma patients: two parallel randomized, placebo-controlled, double-blind clinical trials. *J Trauma*. 2005;59:8–15; discussion 15–18.
- Rizoli SB, Nascimento B Jr, Osman F, et al. Recombinant activated coagulation factor VII and bleeding trauma patients. *J Trauma*. 2006;61:1419–1425.
- Pusateri AE, Ryan KL, Delgado AV, et al. Effects of increasing doses of activated recombinant factor VII on haemostatic parameters in swine. *Thromb Haemost*. 2005;93:275–283.
- Forgey WW. *The Basic Essentials of Hypothermia*. Guilford, CT: Pequot; 1999.
- Fischer RP, Souba WW, Ford EG. Temperature-associated injuries and syndromes. In: Felician DV, Moore EE, Mattox KL, eds. *Trauma*. 3rd ed. Stamford, CT: Appleton & Lange; 1996:951–952.
- Segel IH. *Enzyme Kinetics: Behavior and Analysis of Rapid Equilibrium and Steady-State Enzyme Systems*. Wiley classics library edition. New York: Wiley; 1993.
- Douning LK, Ramsay MA, Swygert TH, et al. Temperature corrected thrombelastography in hypothermic patients. *Anesth Analg*. 1995;81:608–611.

33. Kettner SC, Kozek SA, Groetzner JP, et al. Effects of hypothermia on thrombelastography in patients undergoing cardiopulmonary bypass. *Br J Anaesth*. 1998;80:313–317.
34. Watts DD, Trask A, Soeken K, Perdue P, Dols S, Kaufmann C. Hypothermic coagulopathy in trauma: effect of varying levels of hypothermia on enzyme speed, platelet function, and fibrinolytic activity. *J Trauma*. 1998;44:846–854.
35. Reed RL II, Johnson TD, Hudson JD, Fischer RP. The disparity between hypothermic coagulopathy and clotting studies. *J Trauma*. 1992;33:465–470.
36. Rohrer MJ, Natale AM. Effect of hypothermia on the coagulation cascade. *Crit Care Med*. 1992;20:1402–1405.
37. Gubler KD, Gentilello LM, Hassantash SA, Maier RV. The impact of hypothermia on dilutional coagulopathy. *J Trauma*. 1994;36:847–851.
38. Sorensen B, Johansen P, Christiansen K, Woelke M, Ingerslev J. Whole blood coagulation thrombelastographic profiles employing minimal tissue factor activation. *J Thromb Haemost*. 2003;1:551–558.
39. Wolberg AS, Meng ZH, Monroe DM III, Hoffman M. A systematic evaluation of the effect of temperature on coagulation enzyme activity and platelet function. *J Trauma*. 2004;56:1221–1228.
40. Martini WZ. The effects of hypothermia on fibrinogen metabolism and coagulation function in swine. *Metabolism*. 2007;56:214–221.
41. Shimokawa M, Kitaguchi K, Kawaguchi M, et al. The influence of induced hypothermia for hemostatic function on temperature-adjusted measurements in rabbits. *Anesth Analg*. 2003;96:1209–1213.
42. Aiyagari V, Menendez JA, Diringer MN. Treatment of severe coagulopathy after gunshot injury to the head using recombinant activated factor VII. *J Crit Care*. 2005;20:176–179.
43. Levy JH, Fingerhut A, Brott T, et al. Recombinant factor VIIa in patients with coagulopathy secondary to anticoagulant therapy, cirrhosis, or severe traumatic injury: review of safety profile. *Transfusion*. 2006;46:919–933.
44. Gerotziakas GT, Chakroun T, Depasse F, Arzoglou P, Samama MM, Elalamy I. The role of platelets and recombinant factor VIIa on thrombin generation, platelet activation and clot formation. *Thromb Haemost*. 2004;91:977–985.
45. Sondeen JL, Pusateri AE, Hedner U, Yantis LD, Holcomb JB. Recombinant factor VIIa increases the pressure at which rebleeding occurs in porcine uncontrolled aortic hemorrhage model. *Shock*. 2004;22:163–168.
46. Lisman T, Adelmeijer J, Cauwenberghs S, Van Pampus EC, Heemskerk JW, De Groot PG. Recombinant factor VIIa enhances platelet adhesion and activation under flow conditions at normal and reduced platelet count. *J Thromb Haemost*. 2005;3:742–751.
47. Butenas S, Brummel KE, Branda RF, Paradis SG, Mann KG. Mechanism of factor VIIa-dependent coagulation in hemophilia blood. *Blood*. 2002;99:923–930.
48. Hoffman M, Monroe DM, Roberts HR. Platelet-dependent action of high-dose factor VIIa. *Blood*. 2002;100:364–365; author reply 365.
49. Veldman A, Hoffman M, Ehrenforth S. New insights into the coagulation system and implications for new therapeutic options with recombinant factor VIIa. *Curr Med Chem*. 2003;10:797–811.
50. Meng ZH, Wolberg AS, Monroe DM III, Hoffman M. The effect of temperature and pH on the activity of factor VIIa: implications for the efficacy of high-dose factor VIIa in hypothermic and acidotic patients. *J Trauma*. 2003;55:886–891.
51. Martinowitz U, Michaelson M. Guidelines for the use of recombinant activated factor VII (rFVIIa) in uncontrolled bleeding: a report by the Israeli Multidisciplinary rFVIIa Task Force. *J Thromb Haemost*. 2005;3:640–648.
52. Arnal D, Pineiro P, Garutti I, Olmedilla L, Sanz J, Lajara A. Recombinant activated factor VII used in a man with refractory bleeding from a stab wound injuring the liver and kidney. *Rev Esp Anesthesiol Reanim*. 2004;51:284–288.
53. Martinowitz U, Holcomb JB, Pusateri AE, et al. Intravenous rFVIIa administered for hemorrhage control in hypothermic coagulopathic swine with grade V liver injuries. *J Trauma*. 2001;50:721–729.
54. Schreiber MA, Holcomb JB, Hedner U, Brundage SI, Macaitis JM, Hoots K. The effect of recombinant factor VIIa on coagulopathic pigs with grade V liver injuries. *J Trauma*. 2002;53:252–257; discussion 257–259.
55. Klemcke HG, Delgado A, Holcomb JB, et al. Effect of recombinant FVIIa in hypothermic, coagulopathic pigs with liver injuries. *J Trauma*. 2005;59:155–161; discussion 161.